



UNIVERSITI PUTRA MALAYSIA

**ISOLATION AND SEQUENCE ANALYSES OF
SALINITY TOLERANCE GENES FROM BRUGUIERA
CYLINDRICA (L.) BLUME**

WONG YEEN YEE

FBSB 2005 17

**ISOLATION AND SEQUENCE ANALYSES OF SALINITY TOLERANCE
GENES FROM *BRUGUIERA CYLINDRICA* (L.) BLUME**

**Disertai CD-ROM / disket yang boleh diperolehi
di Bahagian Media dan Arkib
(*Accompanying CD-ROM / disk available
at the Media and Archives Division*)**

By

WONG YEEN YEE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of
Science**

July 2005



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**ISOLATION AND SEQUENCE ANALYSES OF SALINITY TOLERANCE
GENES FROM *BRUGUIERA CYLINDRICA* (L.) BLUME**

By

WONG YEEN YEE

July 2005

Chairman : Ho Chai Ling, PhD

Faculty : Biotechnology and Biomolecular Sciences

Salinity is a major abiotic stress limiting the productivity of crop plants globally. The discovery of novel genes in stress adaptation will provide effective genetic engineering strategies leading to greater stress tolerance. The objectives of this research are to identify and isolate salinity tolerance genes from the mangrove plant, *Bruguiera cylindrica* (L.) Blume through suppression subtractive hybridization (SSH) and bacterial functional assay.

B. cylindrica propagules were grown in fresh water and 20 ppt salinity water. Root morphology differences between *B. cylindrica* grown in fresh water and 20 ppt salinity water were largely due to the need of roots to obtain more water and nutrients during salinity stress. *B. cylindrica* plants grew better in the presence of salt as higher mean values were obtained for all morphological measurements compared to *B. cylindrica* plants grown in fresh water.

Four RNA extraction methods were attempted to obtain high yield and high purity RNA. The cesium chloride method was chosen for RNA extraction as it gave the highest amount of pure RNA. Subtracted cDNAs were prepared from the roots of the *B. cylindrica* seedlings that were grown in fresh water and salt water, respectively. A total of 84 subtracted cDNAs were cloned into pCR-BLUNT II TOPO and sequenced. A total of 51 subtracted cDNAs with good sequencing quality were assembled into 7 contigs and 10 singletons. These non-redundant sequences were grouped into unknown protein (41.18%), novel (29.41%), protein destination and storage (11.76%), energy (5.88%), intracellular traffic (5.88%) and protein synthesis (5.88%). Some motifs of novel and unknown sequences may involve in the salinity tolerance of *B. cylindrica* such as Kv1.3 voltage-gated K⁺ ion(s) channel signature, calcium-activated BK potassium channel alpha subunit and Kir2.1 inward rectifier K⁺ ion(s) channel signature.

Meanwhile, a cDNA library was also constructed from the roots of *B. cylindrica* that were grown in fresh water. Bacterial functional assay was performed to identify cDNAs that confer salt tolerance. A total of 85 cDNA clones that were able to grow on 2× YT containing 400 mM NaCl were sequenced and 73 cDNAs with good sequence quality were assembled into 9 contigs and 53 singletons. The non-redundant sequences were also categorised into unknown protein (58.06%), metabolism (9.68%), transporters (9.68%), transcription (6.45%), energy (4.84%), cell growth/division (4.84%), novel (3.23%), miscellaneous (1.61%) and disease/defense (1.61%). A motif search on novel and unknown cDNA

sequences had revealed some possible motifs that may be involved in salinity tolerance of *B. cylindrica* e.g. *C. elegans* Srg family integral membrane protein signature and 2Fe-2S ferredoxins, iron-sulfur binding region signature.

Sequence analysis of subtracted cDNAs and putative salt tolerant cDNAs isolated by bacterial functional assay showed some putative proteins that may be involved in the salinity tolerance of *B. cylindrica* such as putative potassium transporter HAK1p (M33), putative zinc finger protein (M3), ubiquitin (BC27) and L-ascorbate peroxidase (A46).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PEMENCILAN DAN ANALISIS TURUTAN GEN-GEN TOLERANSI
KEMASINAN DARI *BRUGUIERA CYLINDRICA* (L.) BLUME**

Oleh

WONG YEEN YEE

Julai 2005

Pengerusi : Ho Chai Ling, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Kemasinan merupakan suatu tekanan abiotik utama yang menghadkan produktiviti tanaman di seluruh dunia. Penemuan gen-gen novel yang terlibat dalam adaptasi tekanan persekitaran akan menyediakan asas strategi pengubahsuaian genetik ke arah penyesuaian tanaman terhadap tekanan persekitaran yang lebih tinggi. Matlamat penyelidikan ini adalah untuk mengenalpasti dan memencilkan gen-gen dalam toleransi kemasinan dari pokok bakau, *Bruguiera cylindrica* (L.) Blume melalui strategi “suppression subtractive hybridization (SSH)” dan “bacterial functional assay”.

Biji benih vivipariti *B. cylindrica* disiram air biasa dan air bergaram sebanyak 20 ppt. Perbezaan morfologi akar yang terhasil di antara *B. cylindrica* yang disiram dengan air biasa dan air bergaram kemungkinan disebabkan oleh keperluan akar untuk memperolehi air dan nutrien yang lebih banyak semasa berada di dalam keadaan tekanan kemasinan. *B. cylindrica* membesar dengan lebih baik dengan kehadiran garam kerana nilai purata

ukuran morfologi yang lebih tinggi diperolehi dibandingkan dengan *B. cylindrica* yang disiram dengan air biasa.

Daripada empat kaedah pengekstrakan RNA yang telah dicuba, kaedah cesium klorida telah dipilih kerana kaedah ini memberi hasil RNA yang paling tulen dan tinggi. cDNA yang diekspres di dalam akar *B. cylindrica* yang membesar dengan kehadiran garam sahaja telah diperolehi melalui strategi SSH. Strategi SSH menyingkirkan cDNA di dalam akar *B. cylindrica* yang disiram dengan air biasa. Sejumlah 81 klon dari SSH telah diklon ke dalam pCR-BLUNT II TOPO dan diujuk. Sebanyak 51 jujukan cDNA yang berkualiti dihimpunkan ke dalam 7 “contig” and 10 “singleton”. Jujukan-jujukan yang tidak redandensi ini dikumpulkan ke dalam kumpulan protin tidak diketahui (41.18%), unik (29.41); destinasi and penyimpanan protin (11.76%), tenaga (5.88%), trafik intrasel (5.88%) dan protin sintesis (5.88%). Motif yang dimiliki oleh jujukan unik dan tidak diketahui mungkin terlibat di dalam toleransi kemasinan pokok bakau *B. cylindrica* seperti Kv1.3 voltage-gated K⁺ ion(s) channel signature, calcium-activated BK potassium channel alpha subunit and Kir2.1 inward rectifier K⁺ ion(s) channel signature.

Selain itu, satu perpustakaan cDNA telah disediakan daripada akar *B. cylindrica* yang disiram dengan air biasa. “Bacterial functional assay” telah digunakan untuk mengenalpasti cDNA yang mempunyai sifat toleransi kemasinan. Sejumlah 85 klon yang tumbuh di atas plat 2× YT telah dipencil dan diujuk. Sebanyak 73 jujukan yang berkualiti dihimpunkan ke dalam 9 “contig” and 53 “singleton”. Jujukan-jujukan yang tidak redandensi turut

dikategorikan ke dalam kumpulan protin tidak diketahui (58.06%), metabolime (9.68%), pengangkutan (9.68%), transkripsi (6.45%), tenaga (4.84%), pertumbuhan and pembahagian sel (4.84%), unik (3.23%), lain-lain (1.61%) serta penyakit dan pertahanan (1.61%). Carian motif ke atas jujukan unik dan tidak diketahui telah memberi gambaran terhadap motif yang mungkin terlibat di dalam toleransi kemasinan *B. cylindrica* seperti *C. elegans* Srg family integral membrane protein signature and 2Fe-2S ferredoxins, iron-sulfur binding region signature.

Analisa penjujukan cDNA dari SSH dan “bacterial functional assay” menunjukkan kehadiran protin-protin yang mungkin terlibat di dalam toleransi kemasinan pokok bakau, *B. cylindrica* seperti “putative potassium transporter HAK1p” (M33), “putative zinc finger protein” (M3), “ubiquitin” (BC27) and “L-ascorbate peroxidase” (A46).

ACKNOWLEDGEMENTS

I would like to express my gratitude and appreciation to Dr. Ho Chai Ling for giving me the opportunity to pursue my studies in the field of molecular biology. Throughout my study period, she provided encouragement, sound advice, good teaching and also for her patience in guiding me. Not to forget Assoc. Prof. Dr Jenni, Assoc. Prof. Dr. Raha and Dr Michael Wong for their guidance and advices throughout my study.

Special thanks to my lab mates: Dr. Sudha, Tony, Seddon, Dang, Swee Sen and many others for their kind assistance, advices and support. I am grateful to Cheong Xin for his help in journals searching; Mr. Lee Weng Wah and Mr. Lee Yang Ping for answering all my queries in sequence analysis.

I would also like to acknowledge the Ministry of Science, Technology and Innovation, Malaysia, for providing financial support of PASCA postgraduate fellowship.

Lastly, and most importantly, I wish to thank my family for their understanding, love and support. To them I dedicate this thesis.

I certify that an Examination Committee met on 25th July 2005 to conduct the final examination of Wong Yeen Yee on her Master of Science thesis entitled "Isolation and Sequence Analyses of Salinity Tolerance Genes from *Bruguiera cylindrica* (L.) Blume" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Norihan Mohd Saleh, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Suhaimi Napis, PhD

Associate Professor
Institute of Multimedia and Software
Universiti Putra Malaysia
(Internal Examiner)

Mohd Puad Abdullah, PhD

Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Helen Nair, PhD

Professor
Faculty of Applied Sciences
Asian Institute of Medicine, Science and Technology (AIMST)
(External Examiner)



GULAM RUSUL RAHMAT ALI, PhD

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: **25 AUG 2005**

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science degree. The members of the Supervisory Committee are as follows:

Ho Chai Ling, PhD

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

Jennifer Ann Harikrishna, PhD

Associate Professor

Malaysia University of Science and Technology

(Member)

Raha Abdul Rahim, PhD

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Member)



AINI IDERIS, PhD

Professor/ Dean


School of Graduate Studies

Universiti Putra Malaysia

Date: **17 NOV 2005**

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



WONG YEEN YEE

Date: 19/10/05

TABLE OF CONTENTS

| | Page |
|---|-----------|
| ABSTRACT | ii |
| ABSTRAK | v |
| ACKNOWLEDGEMENTS | viii |
| APPROVAL | ix |
| DECLARATION | xi |
| LIST OF TABLES | xv |
| LIST OF FIGURES | xvi |
| LIST OF ABBREVIATIONS | xviii |
| CHAPTER | |
| 1 INTRODUCTION | 1 |
| 2 LITERATURE REVIEW | 3 |
| 2.1 Stress in plants | 3 |
| 2.1.1 Salt and drought stress | 3 |
| 2.2 Stress tolerance in plants | 7 |
| 2.2.1 Salt tolerance in plants | 8 |
| 2.3 Mechanism of salt tolerance in plants | 9 |
| 2.3.1 Detoxification | 11 |
| 2.3.2 Homeostasis | 12 |
| 2.3.3 Growth regulation | 22 |
| 2.4 Mangroves | 23 |
| 2.4.1 Importance of mangroves | 25 |
| 2.4.2 <i>Bruguiera cylindrica</i> (L.) Blume | 29 |
| 2.4.3 Salt tolerance in <i>Bruguiera</i> sp. | 33 |
| 2.5 Differentially expressed genes | 36 |
| 2.5.1 Other methods | 39 |
| 2.6 Functional assay | 42 |
| 3 MATERIALS AND METHODS | 46 |
| 3.1 Sample collection and preparation | 46 |
| 3.2 Optimization of RNA extraction | 47 |
| 3.2.1 Method 1: Cesium chloride method (Glišin <i>et al.</i> , 1974) | 47 |
| 3.2.2 Method 2: CTAB (hexacetyltrimethyl ammonium bromide) method (Apt <i>et al.</i> 1995) | 49 |
| 3.2.3 Method 3: CTAB (hexacetyltrimethyl ammonium bromide) method (Apt <i>et al.</i> 1995) with potassium acetate | 50 |
| 3.2.4 Method 4: CTAB (hexacetyltrimethyl ammonium bromide) method (Sugihara <i>et al.</i> 2000) | 50 |
| 3.3 Isolation of mRNA from total RNA | 51 |
| 3.4 Suppression subtractive hybridization (SSH) | 52 |
| 3.4.1 First-strand cDNA synthesis | 52 |



| | | |
|--------|--|----|
| 3.4.2 | Second-strand cDNA synthesis | 52 |
| 3.4.3 | <i>RsaI</i> digestion | 53 |
| 3.4.4 | Adaptor ligation | 54 |
| 3.4.5 | First hybridization | 55 |
| 3.4.6 | Second hybridization | 55 |
| 3.4.7 | Primary PCR amplification | 55 |
| 3.4.8 | Secondary PCR amplification | 56 |
| 3.5 | Analysis of subtracted cDNAs | 56 |
| 3.5.1 | Purification of subtracted cDNAs | 56 |
| 3.5.2 | Preparation of pCR®-Blunt II-TOPO plasmid | 57 |
| 3.5.3 | Ligation of subtracted cDNAs to pCR®-Blunt II-TOPO | 58 |
| 3.5.4 | Transformation of ligated plasmid into TOP10 competent cell | 58 |
| 3.5.5 | Preparation of plasmid DNA | 58 |
| 3.5.6 | Restriction enzymes analysis of transformants | 60 |
| 3.5.7 | PCR analysis of transformants | 60 |
| 3.5.8 | Sequencing | 60 |
| 3.5.9 | Sequence analysis | 61 |
| 3.6 | Yeast functional assay | 62 |
| 3.6.1 | Preparation of pYC2/NT A, B and C | 62 |
| 3.6.2 | Ligation of subtracted cDNAs to pYC2/NT A | 65 |
| 3.6.3 | Transformation of ligated plasmid into TOP10 cells | 65 |
| 3.6.4 | Preparation of plasmid DNA | 65 |
| 3.6.5 | Restriction enzymes analysis of transformants | 65 |
| 3.6.6 | PCR analysis of transformants | 66 |
| 3.6.7 | Sequencing of positive clones | 66 |
| 3.6.8 | Yeast transformation of ligated plasmid into InVSc1 cells | 66 |
| 3.6.9 | PCR confirmation of yeast transformants | 67 |
| 3.6.10 | Salt tolerance testing on yeast transformants | 68 |
| 3.7 | Construction of cDNA library from <i>B. cylindrica</i> roots | 68 |
| 3.7.1 | First-strand cDNA synthesis | 69 |
| 3.7.2 | Second-strand cDNA synthesis | 69 |
| 3.7.3 | Blunting the cDNA termini | 69 |
| 3.7.4 | Ligation of <i>EcoRI</i> adapters | 70 |
| 3.7.5 | Phosphorylating the <i>EcoRI</i> ends | 70 |
| 3.7.6 | <i>XhoI</i> digestion | 71 |
| 3.7.7 | Size fractionation | 71 |
| 3.7.8 | Quantification of cDNA with ethidium bromide | 72 |
| 3.7.9 | Ligation of cDNA to Uni-ZAP XR vector | 72 |
| 3.7.10 | Preparing the host bacteria | 72 |
| 3.7.11 | Packaging the ligated product | 73 |
| 3.7.12 | Primary plating and titrating | 73 |
| 3.7.13 | <i>In vivo</i> excision | 74 |
| 3.7.14 | Amplification of cDNA library | 75 |
| 3.8 | Bacterial functional assay | 76 |
| 3.8.1 | Mass Excision | 76 |
| 3.8.2 | Plating | 77 |

| | | |
|----------|---|------------|
| 4 | RESULTS | 78 |
| 4.1 | Sample collection and preparation | 78 |
| 4.2 | RNA extraction | 81 |
| 4.3 | Isolation of mRNA from total RNA | 84 |
| 4.4 | Suppression subtractive hybridization (SSH) | 84 |
| 4.5 | Yeast functional assay | 92 |
| 4.6 | Bacteria functional assay | 96 |
| 5 | DISCUSSION | 106 |
| 5.1 | Sample collection and preparation | 106 |
| 5.2 | RNA extraction | 110 |
| 5.3 | Suppression subtractive hybridization (SSH) | 111 |
| 5.4 | Yeast functional assay | 115 |
| 5.5 | Bacteria functional assay | 115 |
| 5.6 | The possible salinity tolerance genes in <i>B. cylindrica</i> | 118 |
| 6 | CONCLUSION | 123 |
| 6.1 | Recommended study | 124 |
| | REFERENCES | 126 |
| | APPENDICES | 148 |
| | Appendix A | 148 |
| | Appendix B | 150 |
| | Appendix C | 151 |
| | Appendix D | 152 |
| | Appendix E | 153 |
| | Appendix F | 158 |
| | Appendix G | 162 |
| | Appendix H | 163 |
| | Appendix I | 164 |
| | Appendix J | 172 |
| | Appendix K | 186 |
| | Appendix L | 187 |
| | Appendik M | 188 |
| | BIODATA OF THE AUTHOR | 189 |

LIST OF TABLES

| Table | | Page |
|-------|--|------|
| 1 | Sources of environmental stress for plants (reproduced from Hale and Orcutt, 1987) | 4 |
| 2 | Products of mangrove ecosystem (reproduced from Sanger <i>et al.</i> , 1983) | 28 |
| 3 | Measurements of <i>B. cylindrica</i> plants grown in green house in fresh water and 20 ppt salinity water, respectively | 80 |
| 4 | A comparison of results from four RNA extraction methods | 82 |
| 5 | The assembled contigs of <i>B. cylindrica</i> subtracted cDNAs | 89 |
| 6 | Putative functions of subtracted cDNAs | 91 |
| 7 | Subtracted cDNAs that had been successfully ligated to pYC2/NT | 95 |
| 8 | The assembled contigs from the putative salt tolerant cDNAs of <i>B. cylindrica</i> roots that were isolated by bacterial functional assay | 102 |
| 9 | Putative functions of the salt tolerant cDNAs of <i>B. cylindrica</i> roots that were isolated by bacterial functional assay | 104 |



LIST OF FIGURES

| Figure | | Page |
|--------|--|------|
| 1 | Functional demarcation of salt and drought stress signaling pathways (reproduced from Zhu, 2002). | 10 |
| 2 | Na ⁺ ion(s) uptake and extrusion at the plant plasma membrane (reproduced from Blumwald <i>et al.</i> , 2000). | 17 |
| 3 | Regulation of ion (e.g., Na ⁺ ion(s) and K ⁺ ion(s)) homeostasis by the SOS pathway (reproduced from Zhu, 2003). | 17 |
| 4 | Na ⁺ ion(s) compartmentation at the tonoplast (reproduced from Blumwald <i>et al.</i> , 2000). | 21 |
| 5 | Flowers of <i>B. cylindrica</i> (reproduced from Tan, 2001). | 31 |
| 6 | <i>B. cylindrica</i> plants at Morib, Selangor Darul Ehsan (source: Wong Yeen Yee). | 31 |
| 7 | <i>B. cylindrica</i> roots with kneed pneumatophores (loops protruding out of the sand) (reproduced from Ng and Sivasothi, 2001). | 32 |
| 8 | Young propagule of <i>B. cylindrica</i> at Morib, Selangor Darul Ehsan (source: Wong Yeen Yee). | 32 |
| 9 | Schematic diagram of PCR-Select cDNA subtraction (reproduced from Clontech PCR-Select™ cDNA Subtraction kit user manual, 2002). | 38 |
| 10 | Morphology of 4 month-old <i>B. cylindrica</i> seedlings grew in fresh water and 20 ppt salinity water. | 79 |
| 11 | RNA extracted from <i>B. cylindrica</i> roots by using method 1, method 2, method 3 and method 4. | 83 |
| 12 | Gel electrophoresis of subtracted cDNA library of roots of <i>B. cylindrica</i> grown in fresh water and 20 ppt salinity water. | 86 |
| 13 | Assembly of subtracted cDNAs into contigs using CAP3 sequence assembly program. | 88 |
| 14 | Graphic representation of sequences assembled in contig1. | 90 |
| 15 | The functional classification of 17 non-redundant subtracted cDNAs from the roots of <i>B. cylindrica</i> . | 90 |

| | | |
|----|---|-----|
| 16 | Growth of yeast strain INVSc1 with different OD ₆₀₀ readings on YP induction plates with different concentrations of NaCl. | 94 |
| 17 | Gel electrophoresis of the PCR products of cDNA clones from the roots of <i>B. cylindrica</i> that grew in fresh water. | 97 |
| 18 | Mass-excised cDNA library of <i>B. cylindrica</i> roots plated on 2×YT plate containing 171 mM of NaCl and 400 mM of NaCl. | 97 |
| 19 | Putative salt tolerant colonies that were restreaked on 2× YT (400 mM) for reconfirmation. | 98 |
| 20 | Assembly of cDNAs sequence obtained from bacterial functional assay into contigs using CAP3 sequence assembly program. | 101 |
| 21 | Graphic representation of putative salt tolerant cDNA sequences in contig 6 obtained from bacterial functional assay. | 103 |
| 22 | The functional classification of 62 non-redundant putative salt tolerant cDNA clones of <i>B. cylindrica</i> roots. | 103 |
| 23 | Root morphology of <i>Arabidopsis</i> in high and low concentration of phosphorus, respectively; compared with root morphology of <i>B. cylindrica</i> grown in fresh water and 20 ppt salinity water, respectively | 109 |

LIST OF ABBREVIATIONS

| | |
|-------------|-------------------------------------|
| β | beta |
| λ | lambda |
| $\times g$ | gravitational acceleration |
| μg | microgram |
| μL | microliter |
| $^{\circ}C$ | degree centigrade |
| % | percentage |
| AMV | avian myeloblastosis virus |
| BLAST | Basic Local Alignment Search Tool |
| bp | base pairs |
| BSA | bovine serum albumin |
| Ca | calcium |
| cDNA | complementary DNA |
| CIP | calf intestinal phosphatase |
| Cl | chloride |
| cm | centimeter |
| CsCl | cesium chloride |
| CTAB | hexacetyltrimethyl ammonium bromide |
| dATP | 2'-deoxy-adenosine-5'-triphosphate |
| dCTP | 2'-deoxy-cytidine-5'-triphosphate |
| DEPC | diethyl pyrocarbonate |
| dGTP | 2'-deoxy-guanosine-5'-triphosphate |
| DNA | deoxyribonucleic acid |

| | |
|-------------------|---|
| DNase | deoxyribonuclease |
| dNTPs | deoxynucleotides |
| ds | double-stranded |
| DTT | dithiothreitol |
| dTTP | thymidine-5'-triphosphate |
| EDTA | ethylenediaminetetraacetic acid |
| EtBr | ethidium bromide |
| g | gram |
| HCl | hydrochloric acid |
| K | potassium |
| kb | kilo base pairs |
| L | liter |
| LB | Luria-bertani |
| LiCl | lithium chloride |
| M | molar |
| Mg | magnesium |
| MgSO ₄ | magnesium sulfate |
| mL | milliliter |
| mM | millimolar |
| mRNA | messenger RNA |
| Na | sodium |
| NaCl | sodium chloride |
| NaOAc | sodium acetate |
| NaOH | sodium hydroxide |
| NCBI | National Center for Biotechnology Information |

| | |
|---------------------|---------------------------------------|
| ng | nanogram |
| NH ₄ Oac | ammonium acetate |
| OD | optical density |
| PCR | polymerase chain reactions |
| <i>pfu</i> | plaque forming units |
| ppt | part per thousand |
| RNA | ribonucleic acid |
| RNase A | ribonuclease A |
| ROS | reactive oxygen species |
| rpm | revolution per minute |
| SC-U | SC minimal medium without uracil |
| SDS | sodium dodecyl sulphate |
| SOS | salt overly sensitive |
| SSH | suppression subtractive hybridization |
| U | units |
| v/v | volume per volume |
| w/v | weight per volume |
| YPD | yeast extract peptone dextrose medium |
| YT | Yeast extract tryptone medium |

CHAPTER 1

INTRODUCTION

Salinity is one of the major abiotic stress limiting plant productivity and growth globally. Salinity imposes osmotic stress and ionic stress to the plants. Irrigation practices in agricultural lands have steadily increased the concentration of salt in the soil (Khan and Duke, 2001). Unfavorable physiochemical environments can cause average losses more than 65% of optimal yields (Boyer, 1982). More efficient and productive agriculture will be possible on salt affected soils if crop plants with improved salinity tolerance can be selected and bred through traditional breeding or genetic engineering.

As halophytes can live under high salinity condition, it is advantageous to identify genes that are involved in the salinity tolerance of these plants to adapt to harsh environment. Mangroves are unique communities along the tropical and sub-tropical coastal regions that are formed by almost fifty unrelated plant species (Banzai *et al.*, 2002a). Mangroves are divided into two groups based on their morphological features of salt management i.e. 'secreters' and 'non-secreters' (Tomlinson, 1986). The 'secreters' possess salt glands or salt hairs to eliminate excess salt from plants while the 'non-secreters' have no such morphological devices (Tomlinson, 1986; Banzai *et al.*, 2002a).

In this study, *Bruguiera cylindrica* (L.) Blume, known locally in Malaysia as "bakau putih", was chosen as a source to study novel salinity tolerance

genes in halophytes. *B. cylindrica* is categorized as one of the 'non-secreters' and it tolerates high salinity through the ultrafiltration system whereby the plant is able to exclude a large proportion of salt from the water it uptakes and selectively absorbs only certain ions (Tomlinson, 1986).

The objectives of this study were to identify genes that are involved in salinity tolerance of *B. cylindrica* using suppression subtractive hybridization (SSH) and bacterial functional assay. By using the SSH method, genes that were expressed only in the root of *B. cylindrica* grown in the presence of NaCl can be identified. Meanwhile, *B. cylindrica* cDNAs that confer salt tolerance to bacteria can be isolated by performing bacterial functional assay.

CHAPTER 2

LITERATURE REVIEW

2.1 Stress in plants

Stress results changes in plant physiology caused by one or more environmental and biological factors (Hale and Orcutt, 1987). Table 1 shows the sources of environmental stress for plants (Hale and Orcutt, 1987). Stresses are divided into two groups i.e. abiotic and biotic stress. Abiotic stress depends on geographical and climatic differentiation such as annual rainfall differences, chilling, heat, drought, salinity, flooding and freezing (Holmberg and Bülow, 1998; Abeysinghe *et al.*, 2000). Whereas, examples of biotic stresses are diseases and pests.

2.1.1 Salt and drought stress

Salinity and drought are the two major environmental factors that reduce plant productivity currently (Serrano *et al.*, 1999). Boyer (1982) mentioned that disease and insect damages cause losses of less than 10% while unfavorable physiochemical environments cause losses of more than 65%. From historical records, civilizations had never progressed in one locality for more than 1000-2000 years because of the destruction of the resource base of the area (Ashraf, 1994). As a result of poor water management, civilizations have been destroyed by the accumulation of salt on the soils. If there is limited rainfall, salt is not leached out of the soil. Crop plants take in salt through their roots and yields are reduced as the salt concentration increases. Drought and salinity are interconnected because crop production

Table 1. Sources of environmental stress for plants (reproduced from Hale and Orcutt, 1987).

| Physical | Chemical | Biotic |
|-------------|-----------------------|-------------------|
| Drought | Air pollution | Competition |
| Temperature | Allelochemicals | Allelopathy |
| Radiation | (organic) | Lack of symbioses |
| Flooding | Nutrients (inorganic) | Human activities |
| Mechanical | Pesticides | Diseases |
| Electrical | Toxins | Insects |
| Magnetic | Salts | |
| Wind | pH of soil solution | |